

Some North American Birds Obtained in Japan.—Of North American birds which stand recorded in the literature as having been obtained in Japan, there are: The Buff-breasted Sandpiper, *Tringites subruficollis* (Vieill.), once obtained in the Province of Owari (Stejneger, Proc. U. S. Nat. Mus., Vol. XVI, 1893, p. 616); the Short-billed Gull, *Larus canus brachyrhynchus* Richardson, obtained in the Kurile Islands (Dresser, Man. Pal. Bds., p. 830); and Cassin's Auklet, *Ptychoramphus aleuticus* (Pall.), which occurred also in the Kurile Islands (Uchida, "Ni-honchorui Zuzetsu," Vol. I, p. 301). To the writer are known three more cases of North American birds having been met with as stragglers in Japan. They are as follows:

Mareca americana (Gmelin). BALDPATE.—A nearly adult male of this duck was obtained December 4, 1908, in the duck-decoy pond owned by me at Haneda, between Tokyo and Yokohama (Kuroda, Zool. Mag. Tokyo, Vol. XXI, 1909, p. 145). A second example, an adult male, of the same species, was captured at the same pond, January 16, 1918 (Kuroda, "Tori," Vol. II, No. 6, 1918, p. 52).

This remarkable duck seems to occur in Japan only as a rare straggler in migration, mixed in flocks of the European Widgeon (*M. penelope*). It is said that it is occasionally captured in the neighborhood of Tokyo. I have not yet seen a female example of this duck obtained in this country. Dr. Stejneger (Proc. U. S. Nat. Mus., 1887, p. 136) has recorded that a single straggler of the species was picked up dead on a sand-dune in Bering Island, and Dresser (Man. Pal. Bds., 1903, p. 616) has mentioned that the species occurred two or three times in Great Britain.

Nettion crecca carolinense (Gmelin). GREEN-WINGED TEAL.—A fully adult male, with white crescentic bands on each side of the chest, was obtained at Haneda in the same pond mentioned above, February 17, 1916 (Kuroda, Zool. Mag. Tokyo, Vol. XXVIII, 1916, p. 413). This is the only instance known to me of this teal having occurred in Japan. It is probably a rarer straggler in this country than the American Widgeon. According to Dresser (Man. Pal. Bds., 1903, p. 612) the teal in question had been obtained twice at least in Great Britain.

Xema sabini (J. Sabine). SABINE'S GULL.—A male of this gull in complete summer plumage was collected on the coast of Kesen-numa Bay in Prov. Rikuzen, November, 1909 (Kuroda, Zool. Mag. Tokyo, Vol. XXIV, 1912, p. 55). This is the only case known of its occurrence in Japan.

This Gull like the two species of duck already mentioned is properly an American species, of which some individuals are known to have been met with as accidental stragglers in Europe. Thayer and Bangs (Proc. N. Eng. Zoöl. Club, Vol. V, 1914, p. 11) have noted that it was not observed on the Arctic coast of Siberia, while Koren says he has examined a skin of the species in possession of a native at Nijni Kolymsk, East Siberia.—NAGAMICHI KURODA, *Tokyo, Japan.*